

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO DIVIDEND EXTREME™

EPA REG. NO 100-1141 EPA SLN NO. ID-030019

Seed Treatment That Controls Or Suppresses Certain Seed-Borne And Soil-Borne Diseases In Spring Barley

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions on this Supplemental label and the main EPA-registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This labeling and the federal label for this product must be in the possession of the user at the time of pesticide application.

DO NOT use on agricultural establishments in hopper-box, planter-box, slurry-box, or other seed-treatment applications at or immediately before planting.

Mixing Procedures

Apply Dividend Extreme as a water-based slurry through standard slurry or mist type commercial seed treaters. Prepare a slurry by mixing the appropriate amount of Dividend Extreme in up to 16 fl. oz. of water per 100 lbs. of seed. Mix the slurry thoroughly with the seed to provide uniform coverage.

Spring Barley

Apply Dividend Extreme to barley seed at the rates given in the following table:

Rate/CWT	Diseases Controlled	Diseases Partially Controlled ¹
2 – 4 fl. oz.	Barley Stripe General Seed Rots ² Fusarium Seed Scab Covered Smut Pythium Damping-Off	Take-All Common Root Rot (<i>Cochliobolus spp.</i>) Fusarium Root Rot Fusarium Crown Rot Rhizoctonia Root Rot

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

Important: (1) **DO NOT** use treated seed for feed, food or oil. (2) **DO NOT** graze, cut for green chop silage or hay or otherwise utilize straw or seed screenings as animal feeds from field in which seeds were planted. (3) **DO NOT** plant any crop other than those on this label within 30 days to fields in which treated seeds were planted.

Note: Federal law requires that bags containing treated seeds shall be labeled with the following information: "This seed has been treated with difenconazole fungicide. Do not use for feed, food, or oil purposes. Store away from feeds and foodstuffs." Do not plant any crop other than those on this label within 30 days in fields in which treated seeds were planted.

EPA SLN No: ID-030019

Dividend Extreme™ is a trademark of Syngenta

24(c) Registrant: Syngenta Crop Protection, Inc. Greensboro, NC 27419

Label Code: ID1141059AA0104

Word: IDDividendExtremeBarleySLNLbl 1-15-04

Page 2 of 2 EPA SLN No. ID-030019

²General seed rots controlled include those caused by saprophytic organisms, such as Penicillium and Aspergillus.